

Possible Evidence for a Chiral Particle D_1^X , Axial-Vector in the D Meson System

Daiki ITO, Muneyuki ISHIDA*, Shin ISHIDA, Toshihiko KOMADA**
Hiroshi TONOOKA and Kenji YAMADA**

*Research Institute of Quantum Science, College of Science and Technology
Nihon University, Tokyo 101-0062, Japan*

**Department of Physics, Tokyo Institute of Technology, Tokyo 152-8551, Japan*

***Department of Engineering Science, Junior College Funabashi Campus
Nihon University, Funabashi 274-8501, Japan*

In the covariant level-classification scheme of hadrons proposed recently, the existence of “chiral particles”, scalar and axial-vector mesons as the partners of the ground-state pseudo-scalar and vector mesons, respectively, were predicted in the D and B meson systems, realizing a linear representation of chiral symmetry. In this work we reanalyze the $D^*\pi$ mass spectra obtained through the $\Upsilon(4S)$ and Z^0 decays to show a possibility for the D_1^X .

§1. Introduction

The level classification of hadrons had been done rather successfully on the basis of the non-relativistic quark model(NRQM) for these 4 decades. However, this non-relativistic scheme seems now to be confronted with a serious difficulty: Recently existence^{1), 2)} of the light-mass iso-scalar scalar meson to be identified with σ ^{3), 4)}, the chiral partner of the π meson as a Nambu-Goldstone boson, has been accepted widely⁵⁾. Successively possible existence of the $\kappa(900)$ meson⁶⁾, iso-spinor scalar meson, has been pointed out. Then we are able to identify naturally the σ -meson nonet⁷⁾ with the members $\{\sigma(600), \kappa(900), a_0(980), f_0(980)\}$.

The mass values ($\leq 1\text{GeV}$) of this scalar nonet are in the region of $q\bar{q}$ ground states in NRQM, while there are no seats for scalars in this model. This has aroused recently a hot controversy on what the quark configuration of σ nonet is.

On the other hand we have proposed a new level-classification scheme^{8), 9)}, which has a relativistically covariant framework and in conformity with the approximate chiral symmetry. In this new classification scheme the “chiral states”, which are out of the framework of the NRQM, are expected to exist in the lower mass region, and the σ -nonet is naturally assigned as the $(q\bar{q})$ relativistic S -wave chiral state. In the heavy-light quark meson system it is expected that the approximate chiral symmetry on the light quark is valid, and the new level-classification scheme anticipates existence of the chiral states¹⁰⁾, scalar mesons and axial-vector mesons, in addition to the conventional pseudo-scalar and vector mesons, in the ground states. The purpose of this work is to investigate the possibility of existence of the chiral axial-vector meson D_1^X in the D meson system by re-analyzing some experimental data reported previously.*

* Preliminary results of this work were reported at the conference Hadron 2001 at Protvino.¹¹⁾

§2. Experimental Data and Project for Reanalysis

(*Experimental data to be analyzed*)

As the experimental data^{12), 13)} to be re-analyzed we choose the mass spectra of $D^{*+}\pi^-$ system obtained through the two processes,

$$\text{CLEO II} : e^+ + e^- \rightarrow \Upsilon(4S) \rightarrow D^{*+}\pi^- + \cdots, \quad (2.1)$$

$$\text{DELPHI} : e^+ + e^- \rightarrow Z^0 \rightarrow D^{*+}\pi^- + \cdots, \quad (2.2)$$

which have at least a necessary statistical accuracy deserved to meaningful analysis. We shall take the two kinds of CLEO data on the process Eq. (2.1), CLEO (a) and CLEO (b); the former are the restricted ones so as to stress relatively the conventional axial-vector meson D_1^0 , while the latter are all the relevant data.

As the irrelevant backgrounds, which come from the other than decay processes of the relevant resonant particles, the CLEO group set up the formula

$$B. G. = \alpha(\Delta M)^\beta \times \exp(-\gamma_1(\Delta M) - \gamma_2(\Delta M)^2 - \gamma_3(\Delta M)^3), \quad (2.3)$$

$$\Delta M \equiv M(D^{*}\pi) - (m_{D^{*}} + m_\pi),$$

with fitting parameters $\alpha, \beta, \gamma_1, \gamma_2, \gamma_3$, while the DELPHI group applies the similar formula to Eq. (2.3) with fitting parameters α, β, γ_1 , setting $\gamma_2 = \gamma_3 = 0$. We shall apply the same formula to Eq. (2.3) with parameters $\alpha, \beta, \gamma_1, \gamma_2, \gamma_3$ for the backgrounds, taking into account the possible effects of intermediate D_1^χ production. (*Relevant intermediate resonances*)

The conventional three $(c\bar{q}) - P$ wave mesons may contribute, as intermediate states, to the final $D^{*+}\pi^-$ system with the respective angular momenta $l = 0, 1, 2$ as

$$D_1^* \rightarrow D^{*+} + \pi^- \quad (S - \text{wave}), \quad (2.4)$$

$$D_1 \rightarrow D^{*+} + \pi^- \quad (D - \text{wave}), \quad (2.5)$$

$$D_2^* \rightarrow D^{*+} + \pi^- \quad (D - \text{wave}), \quad (2.6)$$

where D_1^* , D_1 and D_2^* have, respectively, $j_q L_J = 1/2 P_1$, $3/2 P_1$ and $3/2 P_2$ ($j_q = S_q + L$ is the total angular momentum of the light quark), and the respective partial wave states in Eqs. (2.4, 2.5, 2.6) are deduced from the heavy quark symmetry (HQS). In this work, a possible contribution from, in addition to the above conventional resonances, the chiral axial-vector meson D_1^χ is taken into account as

$$D_1^\chi \rightarrow D^{*+} + \pi^- \quad (S - \text{wave}), \quad (2.7)$$

where we inferred that the S -wave decay is dominant because of small Q -value.

Out of the three resonances in Eqs (2.4, 2.5, 2.6) the D_1 and the D_2^* , decaying in the D -wave, have comparatively a small decay width ~ 20 MeV, while D_1^* , decaying in the S -wave, has a large width ~ 300 MeV, reflecting the physical situation that all three resonances have small Q -value and that the width contains a kinematical factor $\Gamma \propto P^{2l+1}$ (P being relative momentum of the final system).

Because of its large width the contribution of D_1^* is difficult to be discriminated from the backgrounds and it was treated as being included into the background in the original analysis by the experimental groups. In this work we shall follow this prescription.

Thus, in order to analyze the relevant mass spectra, we shall take into account the three resonances, D_1 , D_2^* and D_1^χ .

(*Method of analysis*)

We shall apply the VMW method in our relevant case, where the absolute amplitude squared is given by

$$|M(s)|^2 = \sum_i |r_i \Delta_i(s)|^2 + \text{B.G.} , \quad (2.8)$$

$$\Delta_i \equiv \frac{-m_i \Gamma_i}{s - m_i^2 + im_i \Gamma_i} .$$

Here the summation on the suffix i goes over the resonant particles D_1 , D_2^* and D_1^χ , $m_i(\Gamma_i)$ are their masses (widths), and r_i represent their production strength.

The relevant mass spectrum is given by

$$\Gamma(s) = \frac{1}{2\sqrt{s}} \int \frac{d^3 \mathbf{P}_{D^*} d^3 \mathbf{P}_\pi}{(2\pi)^3 2E_{D^*} (2\pi)^3 2E_\pi} \times (2\pi)^4 \delta^{(4)}(P - P_{D^*} - p_\pi) |M(s)|^2 , \quad (2.9)$$

where $s \equiv -(P_\mu)^2$, and P_μ being the total 4-momentum of relevant system.

(*Project for applying χ^2 -analysis*)

In order to obtain the least- χ^2 solution on our relevant parameters, we set up the restrictions on the values of resonance masses and widths as is collected in Table I.

Table I. Restricted regions of values of masses and widths (MeV).

	D_1^χ	D_1	D_2^*
m	2295-2320	2400-2450	2420-2480
Γ	10-60	10-35	10-100

The regions for D_1 and D_2^* are determined properly from the center values and errors of respective resonances reported in PDG table. Concerning a possible candidate for the chiral particle D_1^χ , some excess structure over backgrounds in the regions centered at $m \sim 2300$ MeV, (which is seen commonly through all the three experimental data of CLEO (a) and (b), and of DELPHI) is identified as being due to production of D_1^χ . The restrictions on D_1^χ come simply from inspection of this structure.

We shall perform the two kinds of χ^2 -fitting of the mass spectra, the one with D_1^χ and the other without D_1^χ and compare with their results.

§3. Results of Analysis

The results of our analysis in both the cases, with D_1^χ and without D_1^χ , are shown in Fig. 1: The fitted curves in comparison with the data of CLEO (b), DELPHI

and CLEO (a) are given, respectively, in Fig. 1(a), (b) and (c) [(d), (e) and (f)] in the case with D_1^X [without D_1^X]. The respective contributions from the D_1^X , D_1 , D_2^* and B. G. are also shown.

Actually we have made our analysis in the two steps(, taking into account the difference of statistical accuracy of respective data): First fitting CLEO (b) and DELPHI commonly and secondly fitting CLEO (a) with the fixed values of resonance mass and width obtained through the first fitting.




Fig. 1. Fitted curves in the case with (without) D_1^X and experimental data of CLEO (b) and DELPHI are shown, respectively, in (a) and (c) ((b) and (d)).

The obtained values of mass and width of the relevant resonances are collected in Table II. The values of D_1 and D_2^* are close to those given in PDG table.

In Table III the values of χ^2 and of $\tilde{\chi}^2 (\equiv \chi^2 / \text{No. of data points} - \text{No. of param.})$ in the respective steps of fitting are also given.

Through the inspection of fitted curves in Fig. 1, we may conclude that our results give some evidence for existence of D_1^X , although there is no significant im-




Fig. 2. Fitted curves in the case with (without) D_1^χ are, for reference, given in comparison with experimental data, CLEO(a), where the values of mass and width of relevant resonances are determined in the analysis of CLEO(b) and DELPHI.

Table II. Fitted values of mass and width of resonances (in MeV)

	D_1^χ	D_1	D_2^*
m	2312	2421	2465
Γ	23.03	30.73	42.54

Table III. Values of χ^2 and $\tilde{\chi}^2$

	(A) with D_1^χ			(B) without D_1^χ		
	CLEO (b)	DELPHI	CLEO (a)	CLEO (b)	DELPHI	CLEO (a)
$\tilde{\chi}^2$	$\frac{110}{(140-20)} = 0.923$		$\frac{80.6}{(70-8)} = 1.30$	$\frac{119}{(140-16)} = 0.957$		$\frac{85.5}{(70-7)} = 1.36$
χ^2	59.51	51.28		66.41	52.25	

provement on the reduced χ^2 given in Table III.

§4. Concluding Remarks

We have made reanalysis of the $(D^{*+}\pi^-)$ mass spectra obtained through the processes Eqs. (2.1) and (2.2), by applying the formulas Eqs. (2.8) and (2.9). As the intermediate resonant particles we have taken into account the D_1^χ , a new chiral axial-vector meson outside of the conventional level classification scheme. As a result we have obtained a possible evidence for D_1^χ with $m = 2312$ MeV and $\Gamma = 23.0$ MeV. The obtained values of reduced χ square, $\tilde{\chi}^2 \equiv \chi^2/(\text{No. of data points} - \text{No. of param.})$, are $\tilde{\chi}^2 = 110/(140 - 20) = 0.923$ for the case with D_1^χ and $119/(140 - 16) = 0.957$ for the case without D_1^χ , respectively. However, the statistical accuracy of the relevant data is very poor. It is required to have the more accurate data in order to get a definite conclusion.

A possible evidence for another chiral particle B_0^χ , scalar meson in the B system,

is also shown by reanalysis of the $B\pi$ mass spectra through the decay process of Z^0 -boson. It will be given in a separate paper.¹⁴⁾

Acknowledgement

The authors should like to express their gratitude to Professors, K. Takamatsu and T. Tsuru, for their useful discussions and encouragement.

References

- [1] R. Delbourgo and M. D. Scadron, Phys. Rev. Lett. **48** (1982), 765.
- [2] T. Hatsuda and T. Kunihiro, Prog. Theor. Phys. **74** (1985), 765; Phys. Rep. **247** (1994), 221.
- [3] N. N. Achasov and G. N. Schestakov, Phys. Rev. D **49** (1994), 5779.
R. Kaminski, L. Lesniak and J. -P. Maillet, Phys. Rev. D **50** (1994), 3145.
N. A. Tornqvist, Z. Phys. C **68** (1995), 647.
N. A. Tornqvist and M. Roos, Phys. Rev. Lett. **76** (1996), 1575.
M. Harada, F. Sannino and J. Schechter, Phys. Rev. D **54** (1996), 1991.
- [4] S. Ishida, M. Ishida, H. Takahashi, T. Ishida, K. Takamatsu and T. Tsuru, Prog. Theor. Phys. **95** (1996), 745; **98** (1997), 1005.
- [5] Proceedings on *Possible Existence of σ -Meson and Its Implications to Hadron Physics, YITP, Kyoto, 2000* (“ σ -Meson 2000”), Soryushiron Kenkyu (Kyoto) **102** (2001), E1; KEK-proceedings/2000-4.
- [6] S. Ishida, M. Ishida, T. Ishida, K. Takamatsu and T. Tsuru, Prog. Theor. Phys. **98** (1997), 159.
- [7] M. Ishida, Prog. Theor. Phys. **101** (1999), 661.
- [8] S. Ishida, M. Ishida and T. Maeda, Prog. Theor. Phys. **104** (2000), 785.
S. Ishida, M. Ishida and T. Maeda, in *Proc. 8th Int. Conf. on Hadron Spectroscopy, Beijing, China, 24-28 Aug. 1999* (Hadron 99); hep-ph/0002095.
S. Ishida and M. Ishida, in *Proc. Possible Existence of σ -Meson and Its Implications to Hadron Physics, YITP, Kyoto, 2000* (σ -Workshop), Soryushiron Kenkyu (Kyoto) **102** (2001), E1; KEK-proceedings/2000-4.
- [9] S. Ishida and M. Ishida, Phys. Lett. B **539** (2002), 249.
- [10] M. Ishida and S. Ishida, Prog. Theor. Phys. **106** (2001), 373.
- [11] K. Yamada et al., in *HADRON SPECTROSCOPY, Proc. of the Ninth International Conference on Hadron Spectroscopy, IHEP, Protvino, Russia, 2002*, ed. D. Amelin and A. M. Zaitsev, AIP Conf. Proc. Vol. 619 (AIP, New York, 2002), p. 657; hep-ph/0111110.
- [12] CLEO collaboration, P. Avery et al., Phys. Lett. B **331** (1994), 236.
- [13] DELPHI collaboration, P. Abreu et al., Phys. Lett. B **426** (1998), 231.
- [14] D. Ito, M. Ishida, S. Ishida, T. Komada, T. Maeda and I. Yamauchi, to be published.